



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Borrelia burgdorferi sensu lato infection pressure shapes innate immune gene evolution in natural rodent populations across Europe

Tschirren, Barbara

Abstract: Although parasite-mediated selection is assumed to be the main driver of immune gene evolution, empirical evidence that parasites induce allele frequency changes at host immune genes in time and/or space remains scarce. Here, I show that the frequency of a protective gene variant of the innate immune receptor Toll-like receptor 2 in natural bank vole (*Myodes glareolus*) populations is positively associated with the strength of *Borrelia burgdorferi sensu lato* infection risk across the European continent. Thereby, this study provides rare evidence for the role of spatially variable infection pressures in moulding the vertebrate immune system.

DOI: <https://doi.org/10.1098/rsbl.2015.0263>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-117258>

Journal Article

Accepted Version

Originally published at:

Tschirren, Barbara (2015). *Borrelia burgdorferi sensu lato* infection pressure shapes innate immune gene evolution in natural rodent populations across Europe. *Biology Letters*, 11(5):20150263.

DOI: <https://doi.org/10.1098/rsbl.2015.0263>

1 ***Borrelia burgdorferi* sensu lato infection pressure shapes innate**
2 **immune gene evolution in natural rodent populations across Europe**

3

4

5 Barbara Tschirren

6

7 Institute of Evolutionary Biology and Environmental Studies, University of
8 Zurich, Zurich, Switzerland

9

10 Contact Information: barbara.tschirren@ieu.uzh.ch

11

12 **Running title:** *Borrelia* shapes *TLR2* evolution

13

14 Abstract

15 Although parasite-mediated selection is assumed to be the main driver of
16 immune gene evolution, empirical evidence that parasites induce allele
17 frequency changes at host immune genes in time and / or space remains
18 scarce. Here I show that the frequency of a protective gene variant of the
19 innate immune receptor Toll-like receptor 2 (*TLR2*) in natural bank vole
20 (*Myodes glareolus*) populations is positively associated with the strength of
21 *Borrelia burgdorferi* sensu lato infection risk across the European continent.
22 Thereby, this study provides rare evidence for the role of spatially variable
23 infection pressures in moulding the vertebrate immune system.

24

25 Keywords: host-parasite interactions; Lyme disease; wildlife disease; genetic
26 variation; resistance evolution; rodents

27

28 1. Introduction

29 Parasites negatively affect host fitness and thereby select for host responses
30 that prevent or reduce infection. The immune system has a key role in host
31 defence, and parasites are therefore assumed to be main drivers of immune
32 gene evolution [1]. Associations between immune gene variants and parasite
33 resistance are regularly observed in natural populations [2-4]. However, in
34 most cases it is difficult to track changes in allele frequencies over time to
35 directly demonstrate parasite-driven evolution of the hosts' immune system in
36 the wild [5].

37 As an alternative, spatial variation in parasite abundance might provide
38 insights into the role of infection pressures in shaping immune system
39 evolution. In humans, for example, spatial variation in *Plasmodium* sp.
40 prevalence is a strong predictor of the frequency of malaria resistance alleles
41 across countries [6-8]. Evidence for an association between parasite pressure
42 and the frequency of resistance alleles in populations of non-human
43 vertebrates, however, remains scant because of a lack of information on
44 spatial variation in the prevalence of wildlife disease, but also, or mainly, the
45 scarcity of candidate immune genes [9].

46 *Borrelia burgdorferi* sensu lato is a common tick-transmitted pathogen in
47 rodents [10], and the causative agent of human Lyme borreliosis (LB), the
48 most common vector-borne disease in Europe and North America [11, 12].
49 The innate immune receptor Toll-like receptor 2 (*TLR2*) plays a key role in the
50 recognition of bacterial lipoproteins [13] and has been identified as a
51 candidate gene for *Borrelia* resistance in laboratory mice [14, 15].

Furthermore, bank voles (*Myodes glareolus*) carrying certain *TLR2* variants (*TLR2* c₂ [16]) have a substantially reduced probability of becoming *Borrelia* infected (i.e. *TLR2* c₂ confers partial *Borrelia* resistance), highlighting that *TLR2* mediates host-*Borrelia* interactions also in the wild [16]. Based on this knowledge, I hypothesised that *Borrelia*-mediated selection shapes the evolution of *TLR2* in rodents. Indeed, substantial variation in the frequency of the protective *TLR2* variant has been observed in bank vole populations across Europe [17]. Although, data on *Borrelia* prevalence in these populations are not available, rates of human LB cases are reported for most European countries [18], and can be used as a proxy for *Borrelia* infection risk.

2. Material and methods

Information on Lyme borreliosis (LB) incidence in a country, measured as the average annual number of LB cases per 100'000 inhabitants, was obtained from [18] and [19] (Table 1). Data on the frequency of the protective *TLR2* c₂ variant in the local bank vole population were obtained from [17] (Table 1). Eight well-defined mitochondrial bank vole lineages have been described in Europe [20-22], which reflect the colonisation history of the species after the last glaciation. Phylogeographic analyses based on the mitochondrial cytochrome *b* gene were used to identify to which lineage the different study populations belong [17] (Table 1). For the UK, Germany and Italy two populations were sampled (Table 1). However, the results did not change qualitatively when randomly excluding one of the two populations per country from the analysis.

The frequency of *TLR2* c₂ in the sampled bank vole populations was analysed using a quasibinomial generalised linear model in R [23]. LB incidence in a country was included in the model. In addition, I included the mitochondrial lineage to which a population belongs to account for potential differences in *TLR2* c₂ frequency among populations due to the colonisation history of bank voles in Europe [17, 20-22]. The significance of factors was determined by comparing two nested models, with and without the factor of interest, using likelihood-ratio tests.

3. Results and discussion

As predicted if *Borrelia*-mediated selection affects the large-scale geographical distribution of the protective *TLR2* variant, there was a positive association between the frequency of *TLR2* c₂ in bank vole populations and human LB incidence across 19 European countries ($\chi^2 = 35.636$, $DF = 1$, $P = 0.003$; Fig. 1). No indication was found that *TLR2* c₂ frequency differed among mitochondrial bank vole lineages ($\chi^2 = 13.958$, $DF = 4$, $P = 0.499$), showing that the observed pattern was not influenced by the colonisation history of the species in Europe [17, 20-22]. Rather, the strong association between human LB incidence and the frequency of the protective *TLR2* c₂ variant in bank vole populations indicates that *Borrelia*-mediated selection has shaped the evolution of this innate immune receptor in wild rodents.

Clearly, the average annual rate of human LB cases in a country is only a crude estimate of the actual *Borrelia* exposure bank voles experience at a local scale. Lyme borreliosis incidence rates, for example, show regional differences within countries [18]. Also, there might be differences among

countries in how accurately LB cases are diagnosed and / or reported [18]. Furthermore, lifestyle differences across countries might change the relationship between *Borrelia* abundance and human LB incidence [18]. Finally, human LB cases are partly caused by *Borrelia* genospecies that do not infect rodents (e.g. the bird specialist *Borrelia garinii* [24]). However, all these factors will add noise to the data and therefore reduce the statistical power to detect a relationship between human LB incidence and the frequency of the protective *TLR2* variant in the local bank vole population, rather than generate a spurious correlation. The observed relationship thus likely represents an underestimation of the impact of *Borrelia* pressure on *TLR2* evolution in bank voles.

The finding that the protective *TLR2* c₂ variant occurred at low frequencies in regions where *Borrelia* is rare suggests that resistance may be associated with costs in the absence of the pathogen [25-27]. Indeed, it has been shown that a human Toll-like receptor 4 (*TLR4*) variant reduces mortality during malaria infection, but is disadvantageous in the absence of *Plasmodium* because it increases the susceptibility to severe bacterial infections [7]. Similar trade-offs may maintain the pronounced balanced polymorphism observed at the bank vole *TLR2* [17]. Assessing the costs of carrying resistance alleles in the absence of infection will thus be a fruitful next step to understand the maintenance of immunogenetic variation in wild vertebrates.

In conclusion, this study provides empirical evidence for an association between *Borrelia* infection risk and the frequency of a protective *TLR2* variant in bank vole populations across the European continent. Thereby it is one of

the first to reveal an association between large-scale spatial variation in pathogen pressure and the immunogenetic composition of populations of a wild vertebrate.

Data accessibility

All data supporting this article are presented in Table 1.

Funding statement

The study was financially supported by the University of Zurich Research Priority Program 'Evolution in Action: From Genomes to Ecosystems' and the Swiss National Science Foundation (PP00P3_128386 and PP00P3_157455).

Conflict of interests

I have no competing interests.

Acknowledgements

I thank Erik Postma and two anonymous reviewers for comments on the manuscript. S. Karger AG approved the use of published LB incidence data for this study.

147 References

- 148 [1] Murphy K, Travers P, Walport M. 2008 *Janeway's immunobiology*. 7th ed.
149 ed New York: Garland Science.
- 150 [2] Hill AVS, Allsopp CEM, Kwiatkowski D, Anstey NM, Twumasi P, Rowe PA,
151 Bennett S, Brewster D, McMichael AJ, Greenwood BM. 1991 Common West
152 African HLA antigens are associated with protection from severe malaria.
153 *Nature* **352**, 595-600.
- 154 [3] Bellamy R. 2004 *Susceptibility to Infectious Diseases: The Importance of*
155 *Host Genetics*. New York: Cambridge University Press.
- 156 [4] Piertney SB, Oliver MK. 2006 The evolutionary ecology of the major
157 histocompatibility complex. *Heredity* **96**, 7-21. (doi:10.1038/sj.hdy.6800724).
- 158 [5] Barrett RDH, Schluter D. 2008 Adaptation from standing genetic variation.
159 *Trends Ecol. Evol.* **23**, 38-44. (doi:10.1016/j.tree.2007.09.008).
- 160 [6] Garamszegi LZ. 2014 Global distribution of malaria-resistant MHC-HLA
161 alleles: the number and frequencies of alleles and malaria risk. *Malar. J.* **13**,
162 349. (doi:10.1186/1475-2875-13-349).
- 163 [7] Ferwerda B, McCall MBB, Alonso S, Mouktaroudi M, Giamarellos-
164 Bourboulis EJ, Izagirre N, Syafruddin D, Kibiki G, Cristea T, Hijmans A, et al.
165 2007 TLR4 polymorphisms, infectious diseases, and evolutionary pressure
166 during migration of modern humans. *Proc. Natl. Acad. Sci. USA* **104**, 16645-
167 16650. (doi:10.1073/pnas.0704828104).
- 168 [8] Kwiatkowski DP. 2005 How malaria has affected the human genome and
169 what human genetics can teach us about malaria. *Am. J. Hum. Genet.* **77**,
170 171-192. (doi:10.1086/432519).

171 [9] Spurgin LG, Richardson DS. 2010 How pathogens drive genetic diversity:
 172 MHC, mechanisms and misunderstandings. *Proc. R. Soc. B* **277**, 979-988.
 173 (doi:10.1098/rspb.2009.2084).

174 [10] Kurtenbach K, Hanincova K, Tsao JI, Margos G, Fish D, Ogden NH. 2006
 175 Fundamental processes in the evolutionary ecology of Lyme borreliosis.
 176 *Nature Rev. Microbiol.* **4**, 660-669.

177 [11] Dennis DT, Hayes EB. 2002 Epidemiology of Lyme Borreliosis. In *Lyme*
 178 *Borreliosis: Biology, Epidemiology and Control* (eds Kahl O, Gray JS, Lane
 179 RS, Stanek G), pp. 251-280. Oxford: CABI Publications.

180 [12] Ostfeld RS. 2011 *Lyme Disease: The Ecology of a Complex System*. New
 181 York: Oxford University Press.

182 [13] Medzhitov R. 2001 Toll-like receptors and innate immunity. *Nat. Rev.*
 183 *Immunol.* **1**, 135-145.

184 [14] Wooten RM, Ma Y, Yoder RA, Brown JP, Weis JH, Zachary JF,
 185 Kirschning CJ, Weis JJ. 2002 Toll-like receptor 2 is required for innate, but not
 186 acquired, host defense to *Borrelia burgdorferi*. *J. Immunol.* **168**, 348-355.

187 [15] Roper RJ, Weis JJ, McCracken BA, Green CB, Ma Y, Weber KS,
 188 Fairbairn D, Butterfield RJ, Potter MR, Zachary JF, et al. 2001 Genetic control
 189 of susceptibility to experimental Lyme arthritis is polygenic and exhibits
 190 consistent linkage to multiple loci on chromosome 5 in four independent
 191 mouse crosses. *Genes Immun.* **2**, 388-397.

192 [16] Tschirren B, Andersson A, Scherman K, Westerdahl H, Mittl P, Råberg L.
 193 2013 Polymorphisms at the innate immune receptor *TLR2* are associated with
 194 *Borrelia* infection in a wild rodent population *Proc. R. Soc. B* **280**,
 195 10.1098/rspb.2013.0364.

196 [17] Morger J, Råberg L, Hille SM, Helsen S, Štefka J, Al-Sabi MM, Kapel
 197 CMO, Mappes T, Essbauer S, Ulrich RG, et al. in press Distinct haplotype
 198 structure at the innate immune receptor Toll-like receptor 2 (*TLR2*) across
 199 bank vole populations and lineages in Europe. *Biol. J. Linn. Soc.*

200 [18] Hubálek Z. 2009 Epidemiology of Lyme Borreliosis. In *Lyme Borreliosis:*
 201 *Biological and Clinical Aspects* (eds Lipsker D, Jaulhac B), pp. 31-50. Basel:
 202 Karger.

203 [19] Biletska H, Podavalenko L, Semenyshyn O, Lozynskyj I, Tarasyuk O.
 204 2008 Study of Lyme borreliosis in Ukraine. *Int. J. Med. Microbiol.* **298**, 154-
 205 160. (doi:10.1016/j.ijmm.2008.04.004).

206 [20] Deffontaine V, Libois R, Kotlík P, Sommer R, Nieberding C, Paradis E,
 207 Searle JB, Michaux J. 2005 Beyond the Mediterranean peninsulas: evidence
 208 of central European glacial refugia for a temperate forest mammal species,
 209 the bank vole (*Clethrionomys glareolus*). *Mol. Ecol.* **14**, 1727-1739.
 210 (doi:10.1111/j.1365-294X.2005.02506.x).

211 [21] Kotlík P, Deffontaine V, Mascheretti S, Zima J, Michaux JR, Searle JB.
 212 2006 A northern glacial refugium for bank voles (*Clethrionomys glareolus*).
 213 *Proc. Natl. Acad. Sci. USA* **103**, 14860-14864.
 214 (doi:10.1073/pnas.0603237103).

215 [22] Deffontaine V, Ledevin R, Fontaine MC, Quere JP, Renaud S, Libois R,
 216 Michaux JR. 2009 A relict bank vole lineage highlights the biogeographic
 217 history of the Pyrenean region in Europe. *Mol. Ecol.* **18**, 2489-2502.
 218 (doi:10.1111/j.1365-294X.2009.04162.x).

219 [23] R Development Core Team. 2011 R: A language and environment for
 220 statistical computing. <http://www.R-project.org>.

221 [24] Baranton G, De Martino SJ. 2009 *Borrelia burgdorferi* sensu lato diversity
 222 and its influence on pathogenicity in humans. In *Lyme Borreliosis: Biological
 223 and Clinical Aspects* (eds Lipsker D, Jaulhac B), pp. 1-17. Basel: Karger.
 224 [25] Tschirren B, Richner H. 2006 Parasites shape the optimal investment in
 225 immunity. *Proc. R. Soc. B* **273**, 1773-1777.
 226 [26] Netea MG, Wijmenga C, O'Neill LAJ. 2012 Genetic variation in Toll-like
 227 receptors and disease susceptibility. *Nat. Immunol.* **13**, 535-542.
 228 (doi:10.1038/ni.2284).
 229 [27] Sheldon BC, Verhulst S. 1996 Ecological immunology: Costly parasite
 230 defences and trade-offs in evolutionary ecology. *Trends Ecol. Evol.* **11**, 317-
 231 321.
 232
 233

234
 235 Table 1 Frequency of the protective *TLR2* c_2 variant (*TLR2* c_2) in bank vole
 236 populations across Europe and average number of annual Lyme borreliosis
 237 cases per 100'000 inhabitants (*LB incidence*) in the same country. *LB*
 238 incidence data were obtained from [18] and [19]. *N*: number of sequenced
 239 bank voles; *Lineage*: mitochondrial lineage of the local bank vole population.
 240 The exact bank vole sampling locations and *TLR2* and *cytb* genotyping
 241 procedures are described in [17].

242


Country	<i>N</i>	<i>TLR2</i>c_2	<i>Lineage</i>	<i>LB incidence</i>
Austria	17	0.27	Western	130
Belgium	19	0.05	Western	12.58
Czech Republic	19	0.00	Western	31.73
Denmark	19	0.05	Eastern	1.68
England	17	0.00	Western	1.72
Finland	20	0.03	Eastern	18.46
Germany 1	21	0.10	Western	25
Germany 2	18	0.08	Eastern	25
Italy 1	21	0.00	Italian	0.02
Italy 2	19	0.00	Italian	0.02
Lithuania	20	0.18	Carpathian	42.93
Netherlands	10	0.00	Western	2.01
Norway	21	0.00	Carpathian	4.5
Poland	12	0.29	Eastern	9.29
Russia	21	0.07	Eastern	9.24
Scotland	19	0.00	Carpathian	1.72
Slovenia	20	0.30	Western	136.86
Spain	13	0.00	Spanish	9.8
Switzerland	19	0.27	Western	25.09
Sweden	20	0.30	Carpathian	69
Ukraine	20	0.00	Eastern	0.98

243

244 Figure legend

245 Figure 1. Relationship between human Lyme borreliosis incidence (LB cases /
246 100'000 inhabitants) and the frequency of the protective *TLR2* c₂ variant in the
247 local bank vole population across 19 European countries. The circle size
248 indicates the number of sampled *TLR2* alleles (*N*) in a population. Locally
249 fitted smoothed polynomial surface (solid line) and 95% confidence interval
250 (grey area) are presented.

251



252